

Azimuthal Angle Correlations for Rapidity Separated Hadron Pairs in $d + Au$ Collisions at $\sqrt{s_{NN}} = 200$ GeV

S.S. Adler,⁵ S. Afanasiev,²⁰ C. Aidala,¹⁰ N.N. Ajitanand,⁴⁴ Y. Akiba,^{21,40} A. Al-Jamel,³⁵ J. Alexander,⁴⁴ K. Aoki,²⁵ L. Aphecetche,⁴⁶ R. Armendariz,³⁵ S.H. Aronson,⁵ R. Averbek,⁴⁵ T.C. Awes,³⁶ V. Babintsev,¹⁷ A. Baldisseri,¹¹ K.N. Barish,⁶ P.D. Barnes,²⁸ B. Bassalleck,³⁴ S. Bathe,^{6,31} S. Batsouli,¹⁰ V. Baublis,³⁹ F. Bauer,⁶ A. Bazilevsky,^{5,41} S. Belikov,^{19,17} M.T. Bjornndal,¹⁰ J.G. Boissevain,²⁸ H. Borel,¹¹ M.L. Brooks,²⁸ D.S. Brown,³⁵ N. Bruner,³⁴ D. Bucher,³¹ H. Buesching,^{5,31} V. Bumazhnov,¹⁷ G. Bunce,^{5,41} J.M. Burward-Hoy,^{28,27} S. Butsyk,⁴⁵ X. Camard,⁴⁶ P. Chand,⁴ W.C. Chang,² S. Chernichenko,¹⁷ C.Y. Chi,¹⁰ J. Chiba,²¹ M. Chiu,¹⁰ I.J. Choi,⁵³ R.K. Choudhury,⁴ T. Chujo,⁵ V. Cianciolo,³⁶ Y. Cobigo,¹¹ B.A. Cole,¹⁰ M.P. Comets,³⁷ P. Constantin,¹⁹ M. Csanád,¹³ T. Csörgő,²² J.P. Cussonneau,⁴⁶ D. d'Enterria,¹⁰ K. Das,¹⁴ G. David,⁵ F. Deák,¹³ H. Delagrangé,⁴⁶ A. Denisov,¹⁷ A. Deshpande,⁴¹ E.J. Desmond,⁵ A. Devismes,⁴⁵ O. Dietzsch,⁴² J.L. Drachenberg,¹ O. Drapier,²⁶ A. Drees,⁴⁵ A. Durum,¹⁷ D. Dutta,⁴ V. Dzhordzhadze,⁴⁷ Y.V. Efremenko,³⁶ H. En'yo,^{40,41} B. Espagnon,³⁷ S. Esumi,⁴⁹ D.E. Fields,^{34,41} C. Finck,⁴⁶ F. Fleuret,²⁶ S.L. Fokin,²⁴ B.D. Fox,⁴¹ Z. Fraenkel,⁵² J.E. Frantz,¹⁰ A. Franz,⁵ A.D. Frawley,¹⁴ Y. Fukao,^{25,40,41} S.-Y. Fung,⁶ S. Gadrat,²⁹ M. Germain,⁴⁶ A. Glenn,⁴⁷ M. Gonin,²⁶ J. Gosset,¹¹ Y. Goto,^{40,41} R. Granier de Cassagnac,²⁶ N. Grau,¹⁹ S.V. Greene,⁵⁰ M. Grosse Perdekamp,^{18,41} H.-Å. Gustafsson,³⁰ T. Hachiya,¹⁶ J.S. Haggerty,⁵ H. Hamagaki,⁸ A.G. Hansen,²⁸ E.P. Hartouni,²⁷ M. Harvey,⁵ K. Hasuko,⁴⁰ R. Hayano,⁸ X. He,¹⁵ M. Heffner,²⁷ T.K. Hemmick,⁴⁵ J.M. Heuser,⁴⁰ P. Hidas,²² H. Hiejima,¹⁸ J.C. Hill,¹⁹ R. Hobbs,³⁴ W. Holzmann,⁴⁴ K. Homma,¹⁶ B. Hong,²³ A. Hoover,³⁵ T. Horaguchi,^{40,41,48} T. Ichihara,^{40,41} V.V. Ikonnikov,²⁴ K. Imai,^{25,40} M. Inaba,⁴⁹ M. Inuzuka,⁸ D. Isenhower,¹ L. Isenhower,¹ M. Ishihara,⁴⁰ M. Issah,⁴⁴ A. Isupov,²⁰ B.V. Jacak,⁴⁵ J. Jia,⁴⁵ O. Jinnouchi,^{40,41} B.M. Johnson,⁵ S.C. Johnson,²⁷ K.S. Joo,³² D. Jouan,³⁷ F. Kajihara,⁸ S. Kametani,^{8,51} N. Kamihara,^{40,48} M. Kaneta,⁴¹ J.H. Kang,⁵³ K. Katou,⁵¹ T. Kawabata,⁸ A.V. Kazantsev,²⁴ S. Kelly,^{9,10} B. Khachaturov,⁵² A. Khanzadeev,³⁹ J. Kikuchi,⁵¹ D.J. Kim,⁵³ E. Kim,⁴³ G.-B. Kim,²⁶ H.J. Kim,⁵³ E. Kinney,⁹ A. Kiss,¹³ E. Kistenev,⁵ A. Kiyomichi,⁴⁰ C. Klein-Boesing,³¹ H. Kobayashi,⁴¹ L. Kochenda,³⁹ V. Kochetkov,¹⁷ R. Kohara,¹⁶ B. Komkov,³⁹ M. Konno,⁴⁹ D. Kotchetkov,⁶ A. Kozlov,⁵² P.J. Kroon,⁵ C.H. Kuberg,^{1,*} G.J. Kunde,²⁸ K. Kurita,⁴⁰ M.J. Kweon,²³ Y. Kwon,⁵³ G.S. Kyle,³⁵ R. Lacey,⁴⁴ J.G. Lajoie,¹⁹ Y. Le Bornec,³⁷ A. Lebedev,^{19,24} S. Leckey,⁴⁵ D.M. Lee,²⁸ M.J. Leitch,²⁸ M.A.L. Leite,⁴² X.H. Li,⁶ H. Lim,⁴³ A. Litvinenko,²⁰ M.X. Liu,²⁸ C.F. Maguire,⁵⁰ Y.I. Makdisi,⁵ A. Malakhov,²⁰ V.I. Manko,²⁴ Y. Mao,^{38,40} G. Martinez,⁴⁶ H. Masui,⁴⁹ F. Matathias,⁴⁵ T. Matsumoto,^{8,51} M.C. McCain,¹ P.L. McGaughey,²⁸ Y. Miake,⁴⁹ T.E. Miller,⁵⁰ A. Milov,⁴⁵ S. Mioduszewski,⁵ G.C. Mishra,¹⁵ J.T. Mitchell,⁵ A.K. Mohanty,⁴ D.P. Morrison,⁵ J.M. Moss,²⁸ D. Mukhopadhyay,⁵² M. Muniruzzaman,⁶ S. Nagamiya,²¹ J.L. Nagle,^{9,10} T. Nakamura,¹⁶ J. Newby,⁴⁷ A.S. Nyanin,²⁴ J. Nystrand,³⁰ E. O'Brien,⁵ C.A. Ogilvie,¹⁹ H. Ohnishi,⁴⁰ I.D. Ojha,^{3,50} H. Okada,^{25,40} K. Okada,^{40,41} A. Oskarsson,³⁰ I. Otterlund,³⁰ K. Oyama,⁸ K. Ozawa,⁸ D. Pal,⁵² A.P.T. Palounek,²⁸ V. Pantuev,⁴⁵ V. Papavassiliou,³⁵ J. Park,⁴³ W.J. Park,²³ S.F. Pate,³⁵ H. Pei,¹⁹ V. Penev,²⁰ J.-C. Peng,¹⁸ H. Pereira,¹¹ V. Peresedov,²⁰ A. Pierson,³⁴ C. Pinkenburg,⁵ R.P. Pisani,⁵ M.L. Purschke,⁵ A.K. Purwar,⁴⁵ J.M. Qualls,¹ J. Rak,¹⁹ I. Ravinovich,⁵² K.F. Read,^{36,47} M. Reuter,⁴⁵ K. Reygers,³¹ V. Riabov,³⁹ Y. Riabov,³⁹ G. Roche,²⁹ A. Romana,^{26,*} M. Rosati,¹⁹ S.S.E. Rosendahl,³⁰ P. Rosnet,²⁹ V.L. Rykov,⁴⁰ S.S. Ryu,⁵³ N. Saito,^{25,40,41} T. Sakaguchi,^{8,51} S. Sakai,⁴⁹ V. Samsonov,³⁹ L. Sanfratello,³⁴ R. Santo,³¹ H.D. Sato,^{25,40} S. Sato,^{5,49} S. Sawada,²¹ Y. Schutz,⁴⁶ V. Semenov,¹⁷ R. Seto,⁶ T.K. Shea,⁵ I. Shein,¹⁷ T.-A. Shibata,^{40,48} K. Shigaki,¹⁶ M. Shimomura,⁴⁹ A. Sickles,⁴⁵ C.L. Silva,⁴² D. Silvermyr,²⁸ K.S. Sim,²³ A. Soldatov,¹⁷ R.A. Soltz,²⁷ W.E. Sondheim,²⁸ S.P. Sorensen,⁴⁷ I.V. Sourikova,⁵ F. Staley,¹¹ P.W. Stankus,³⁶ E. Stenlund,³⁰ M. Stepanov,³⁵ A. Ster,²² S.P. Stoll,⁵ T. Sugitate,¹⁶ J.P. Sullivan,²⁸ S. Takagi,⁴⁹ E.M. Takagui,⁴² A. Taketani,^{40,41} K.H. Tanaka,²¹ Y. Tanaka,³³ K. Tanida,⁴⁰ M.J. Tannenbaum,⁵ A. Taranenko,⁴⁴ P. Tarján,¹² T.L. Thomas,³⁴ M. Togawa,^{25,40} J. Tojo,⁴⁰ H. Torii,^{25,41} R.S. Towell,¹ V.-N. Tram,²⁶ I. Tserruya,⁵² Y. Tsuchimoto,¹⁶ H. Tydesjö,³⁰ N. Tyurin,¹⁷ T.J. Uam,³² H.W. van Hecke,²⁸ J. Velkovska,⁵ M. Velkovsky,⁴⁵ V. Veszprémi,¹² A.A. Vinogradov,²⁴ M.A. Volkov,²⁴ E. Vznuzdaev,³⁹ X.R. Wang,¹⁵ Y. Watanabe,^{40,41} S.N. White,⁵ N. Willis,³⁷ F.K. Wohn,¹⁹ C.L. Woody,⁵ W. Xie,⁶ A. Yanovich,¹⁷ S. Yokkaichi,^{40,41} G.R. Young,³⁶ I.E. Yushmanov,²⁴ W.A. Zajc,^{10,†} C. Zhang,¹⁰ S. Zhou,⁷ J. Zimányi,²² L. Zolin,²⁰ and X. Zong¹⁹

(PHENIX Collaboration)

¹Abilene Christian University, Abilene, TX 79699, U.S.

²Institute of Physics, Academia Sinica, Taipei 11529, Taiwan

³Department of Physics, Banaras Hindu University, Varanasi 221005, India

- ⁴Bhabha Atomic Research Centre, Bombay 400 085, India
⁵Brookhaven National Laboratory, Upton, NY 11973-5000, U.S.
⁶University of California - Riverside, Riverside, CA 92521, U.S.
⁷China Institute of Atomic Energy (CIAE), Beijing, People's Republic of China
⁸Center for Nuclear Study, Graduate School of Science, University of Tokyo, 7-3-1 Hongo, Bunkyo, Tokyo 113-0033, Japan
⁹University of Colorado, Boulder, CO 80309, U.S.
¹⁰Columbia University, New York, NY 10027 and Nevis Laboratories, Irvington, NY 10533, U.S.
¹¹Dapnia, CEA Saclay, F-91191, Gif-sur-Yvette, France
¹²Debrecen University, H-4010 Debrecen, Egyetem tér 1, Hungary
¹³ELTE, Eötvös Loránd University, H - 1117 Budapest, Pázmány P. s. 1/A, Hungary
¹⁴Florida State University, Tallahassee, FL 32306, U.S.
¹⁵Georgia State University, Atlanta, GA 30303, U.S.
¹⁶Hiroshima University, Kagamiyama, Higashi-Hiroshima 739-8526, Japan
¹⁷IHEP Protvino, State Research Center of Russian Federation, Institute for High Energy Physics, Protvino, 142281, Russia
¹⁸University of Illinois at Urbana-Champaign, Urbana, IL 61801, U.S.
¹⁹Iowa State University, Ames, IA 50011, U.S.
²⁰Joint Institute for Nuclear Research, 141980 Dubna, Moscow Region, Russia
²¹KEK, High Energy Accelerator Research Organization, Tsukuba, Ibaraki 305-0801, Japan
²²KFKI Research Institute for Particle and Nuclear Physics of the Hungarian Academy of Sciences (MTA KFKI RMKI), H-1525 Budapest 114, POBox 49, Budapest, Hungary
²³Korea University, Seoul, 136-701, Korea
²⁴Russian Research Center "Kurchatov Institute", Moscow, Russia
²⁵Kyoto University, Kyoto 606-8502, Japan
²⁶Laboratoire Leprince-Ringuet, Ecole Polytechnique, CNRS-IN2P3, Route de Saclay, F-91128, Palaiseau, France
²⁷Lawrence Livermore National Laboratory, Livermore, CA 94550, U.S.
²⁸Los Alamos National Laboratory, Los Alamos, NM 87545, U.S.
²⁹LPC, Université Blaise Pascal, CNRS-IN2P3, Clermont-Fd, 63177 Aubiere Cedex, France
³⁰Department of Physics, Lund University, Box 118, SE-221 00 Lund, Sweden
³¹Institut für Kernphysik, University of Muenster, D-48149 Muenster, Germany
³²Myongji University, Yongin, Kyonggido 449-728, Korea
³³Nagasaki Institute of Applied Science, Nagasaki-shi, Nagasaki 851-0193, Japan
³⁴University of New Mexico, Albuquerque, NM 87131, U.S.
³⁵New Mexico State University, Las Cruces, NM 88003, U.S.
³⁶Oak Ridge National Laboratory, Oak Ridge, TN 37831, U.S.
³⁷IPN-Orsay, Université Paris Sud, CNRS-IN2P3, BP1, F-91406, Orsay, France
³⁸Peking University, Beijing, People's Republic of China
³⁹PNPI, Petersburg Nuclear Physics Institute, Gatchina, Leningrad region, 188300, Russia
⁴⁰RIKEN (The Institute of Physical and Chemical Research), Wako, Saitama 351-0198, JAPAN
⁴¹RIKEN BNL Research Center, Brookhaven National Laboratory, Upton, NY 11973-5000, U.S.
⁴²Universidade de São Paulo, Instituto de Física, Caixa Postal 66318, São Paulo CEP05315-970, Brazil
⁴³System Electronics Laboratory, Seoul National University, Seoul, South Korea
⁴⁴Chemistry Department, Stony Brook University, SUNY, Stony Brook, NY 11794-3400, U.S.
⁴⁵Department of Physics and Astronomy, Stony Brook University, SUNY, Stony Brook, NY 11794, U.S.
⁴⁶SUBATECH (Ecole des Mines de Nantes, CNRS-IN2P3, Université de Nantes) BP 20722 - 44307, Nantes, France
⁴⁷University of Tennessee, Knoxville, TN 37996, U.S.
⁴⁸Department of Physics, Tokyo Institute of Technology, Oh-okayama, Meguro, Tokyo 152-8551, Japan
⁴⁹Institute of Physics, University of Tsukuba, Tsukuba, Ibaraki 305, Japan
⁵⁰Vanderbilt University, Nashville, TN 37235, U.S.
⁵¹Waseda University, Advanced Research Institute for Science and Engineering, 17 Kikui-cho, Shinjuku-ku, Tokyo 162-0044, Japan
⁵²Weizmann Institute, Rehovot 76100, Israel
⁵³Yonsei University, IPAP, Seoul 120-749, Korea

(Dated: February 8, 2008)

We report on two-particle azimuthal angle correlations between charged hadrons at forward/backward (deuteron/gold going direction) rapidity and charged hadrons at mid-rapidity in deuteron-gold ($d + Au$) and proton-proton ($p + p$) collisions at $\sqrt{s_{NN}} = 200$ GeV. Jet structures are observed in the correlations which we quantify in terms of the conditional yield and angular width of away-side partners. The kinematic region studied here samples partons in the gold nucleus carrying nucleon momentum fraction $x \sim 0.1$ to ~ 0.01 . Within this range, we find no x dependence of the jet structure in $d + Au$ collisions.

PACS numbers: 25.75.Dw, 25.75.Gz

Observations in $d + Au$ collisions at $\sqrt{s_{NN}} = 200$ GeV at the Relativistic Heavy Ion Collider (RHIC) reveal a significant suppression of hadron production at forward rapidity (deuteron-going direction) relative to $p + p$ reactions scaled up by the equivalent number of nucleon-nucleon collisions (N_{coll}) [1, 2, 3]. This suppression is observed for hadrons with momentum transverse to the beam direction over the range $p_T \approx 1.5 - 4$ GeV/ c . In contrast, measurements at mid-rapidity [4, 5, 6, 7] and backward rapidity [1, 2, 3] show a modest enhancement of the particle yield relative to N_{coll} scaling over the same p_T range. Particle production at forward rapidity is sensitive to partons in the gold nucleus which carry a small nucleon momentum fraction (small Bjorken x). The suppression has generated significant theoretical interest including different calculational frameworks for understanding the data [8, 9, 10, 11].

One such framework, the Color Glass Condensate (CGC), attempts to describe the data in terms of gluon saturation [8]. At small x the probability of emitting an extra gluon is large and the number of gluons grows in a limited transverse area. When the transverse density becomes large, partons start to overlap and gluon-gluon fusion processes start to dominate the parton evolution in the hadronic wave functions. As a result the gluon density becomes saturated. Since the nonlinear growth of the gluon density depends on the transverse size of the system, the effects of gluon saturation are expected to set in earlier (at higher Bjorken x) for heavy nuclei accelerated at ultra-relativistic energies than for free nucleons.

In the leading order pQCD framework, a quark or gluon jet with large transverse momentum produced in a hard scattering process (high momentum transfer or large Q^2) must be momentum balanced by another quark or gluon jet in the opposite direction but with almost the same transverse momentum. Thus the azimuthal angle correlation between particles from the pair of jets (referred to as di-jets) is characterized by two peak structures separated by 180 degrees. In CGC calculations, the momentum to balance a jet may come from a large multiplicity of gluons in the saturation regime, and thus no single partner jet may appear on the opposite side [12]. This effect is analogous to the nuclear Mössbauer effect, and is often referred to as the appearance of mono-jets. Alternative calculations, describing the suppression of single hadrons at forward rapidity in $d + Au$ reactions in terms of leading twist pQCD effects, predict no such mono-jet feature [13].

We want to probe this high gluon density regime in $d + Au$ collisions with relatively high transverse momentum particles at forward rapidity. Such particles are likely to result from hard-scattering collisions involving small x partons in the gold nucleus. At small x the gluon density increases rapidly with Q^2 and saturation effects may be relevant for $x \approx 0.01$ at modest p_T . CGC calculations [12] predict significant suppression of the condi-

tional yield and widening of away-side jet azimuthal correlations between rapidity-separated hadron pairs when one of those hadrons is at forward rapidity.

In this Letter we report on measurements of two-particle azimuthal angle correlations between unidentified charged hadrons in $p + p$ and $d + Au$ collisions at $\sqrt{s_{NN}} = 200$ GeV. In our analysis, the two particles are referred to as the *trigger* and *associated* particles. The trigger particle is at forward ($1.4 < \eta < 2.0$) or backward ($-2.0 < \eta < -1.4$) rapidity and the associated particle is at mid-rapidity, $|\eta| < 0.35$. The particles are separated by an average pseudorapidity gap $< \Delta\eta > \sim 1.5$. The criteria for trigger particles, associated particles and event selection are described elsewhere [3, 14]. The two-particle azimuthal angle correlation technique has been used extensively by RHIC experiments and is described in detail elsewhere [14, 15, 16, 17, 18]. In this technique the azimuthal correlation function is formed from the angular difference, $\Delta\phi = \phi^{assoc} - \phi^{trig}$, between each trigger and associated particle pair. Two jet peaks are normally observed in such correlation functions: the near-side peak ($\Delta\phi \sim 0$) in which the two particles come from the same jet, and the away-side peak ($\Delta\phi \sim \pi$) in which they come from the back-to-back jets. In addition to these peaks the correlation functions usually also have a $\Delta\phi$ independent combinatoric background contribution which is due to trigger-associated pairs from different jets or from non-jet processes.

We can construct separate correlation functions that are sensitive to partons in the gold nucleus with different Bjorken x ranges. By choosing trigger particles with $1.0 < p_T < 5.0$ GeV/ c at forward (backward) rapidity, we sample partons in gold nuclei with $x \sim 0.01(0.1)$. At $x \sim 0.1$, we do not expect to be sensitive to any saturation effects, but we may be sensitive at $x \sim 0.01$ [19]. The comparison in $d + Au$ reactions between these two cases, as well as with the $p + p$ case, may give insights into possible saturation effects on jet production and other mechanisms for forward rapidity single particle suppression. It should be noted that the prediction of mono-jets in [12] assumes one particle at pseudorapidity $\eta = 3.8$ and one at mid-rapidity, thus demonstrating sensitivity at even lower x_{Au} ($\sim 10^{-4}$) which is outside the range presented in this analysis.

Data for this analysis were collected by the PHENIX experiment [20] in the 2003 RHIC $p + p$ and $d + Au$ running period. In the case of $d + Au$ collisions, we divide the data into two centrality (impact parameter) classes based on the number of hits in the backward-rapidity PHENIX Beam-Beam Counter (BBC, $-3.9 < \eta < -3.0$). Central (peripheral) collisions comprise 0 – 40% (40 – 88%) of the minimum bias cross section respectively. Using a Glauber model [3] and a simulation of the BBC, we determine $< N_{coll} > = 4.7 \pm 0.4$ for peripheral collisions and $< N_{coll} > = 13.0 \pm 0.9$ for central collisions.

The trigger particles are measured in the PHENIX

muon spectrometers [20]. In this analysis we only select trigger particles from $1.4 < |\eta| < 2.0$ to obtain homogenous acceptance in transverse momentum from $1 < p_T < 5$ GeV/c and to reduce beam correlated backgrounds. We identify hadrons, as opposed to muons, in the muon spectrometers by comparing their momentum and penetration depth. This hadron identification method is described elsewhere [3]. It is notable that our trigger hadrons have a modified composition (pion/kaon/proton ratio) relative to that at the collision vertex due to species-dependent nuclear interaction cross sections. Detailed simulations show that kaons make up 65–90% of positively charged trigger particles and pions make up 70–90% of negatively charged trigger particles. The baryon contribution to our trigger particle sample is negligible. We find the two-particle azimuthal angle correlations for positively and negatively charged trigger particles to be consistent and therefore combined the results. The associated particles are unidentified charged hadrons measured in the PHENIX central spectrometers [20] which cover $|\eta| < 0.35$ and in this analysis have $0.5 < p_T < 2.5$ GeV/c. Standard track selection criteria [14] are applied.

For comparison we have also included measurements where trigger particles and associated particles are both measured in the PHENIX central spectrometers at mid-rapidity. The $d + Au$ points for this comparison are from [14] and the $p + p$ point is an extension in p_T of the analysis that was published in [16].

We define the azimuthal angle correlation function as:

$$CF = \frac{dN(\Delta\phi)/d(\Delta\phi)}{acc(\Delta\phi)} \quad (1)$$

where $dN(\Delta\phi)/d(\Delta\phi)$ is the measured two-particle distribution and $acc(\Delta\phi)$ is the two-particle acceptance obtained by an event mixing technique in which we mix trigger particles with associated particles from different events within the same centrality and collision vertex category. This correction is necessary because the PHENIX central arm detector is not azimuthally symmetric and the pair acceptance varies as a function of $\Delta\phi$.

In Figure 1, we show the correlation functions for trigger particles with $p_T = 2 - 5$ GeV/c and associated particles with $p_T = 0.5 - 1.0$ GeV/c. The top panel is for $p + p$ collisions where we have combined the results from forward and backward pseudorapidity since the collision system is symmetric. The middle and bottom panels of Figure 1 show the correlation functions for central $d + Au$ collisions. The middle panel is for the trigger particle at forward rapidity and the bottom panel is for the trigger particle at backward rapidity. A clear peak is seen near $\Delta\phi = \pi$ in all cases corresponding to the away-side jet. It is notable that there is no peak near $\Delta\phi = 0$, as expected, because the rapidity gap between the two particles is larger than the width of the near side jet.




FIG. 1: Azimuthal angle correlation functions. On the plots, the Gaussian widths from the fits and the signal to background ratio integrated over $\pi - 1 < \Delta\phi < \pi + 1$ are shown. Note that the y -axis is zero-suppressed on the middle and bottom panels.

After constructing the correlation functions in various bins in p_T^{assoc} , p_T^{trig} and η^{trig} we used two methods to determine the unnormalized number of trigger-associated particle pairs, N_{pair} , above a constant background. In the first method, we define

$$N_{pair} = \sum_{\Delta\phi=\pi-1}^{\pi+1} CF(\Delta\phi) - \sum_{\Delta\phi=-1}^{+1} CF(\Delta\phi), \quad (2)$$

where the first term is the integral of the correlation function in the area of the correlation peak ($\pi - 1 < \Delta\phi < \pi + 1$) and the second term is the integral away from the peak ($-1 < \Delta\phi < 1$). In the second method we fit the correlation function with a Gaussian distribution centered at $\Delta\phi = \pi$ plus a constant background. The values of N_{pair} obtained by each method are found to be consistent and the small differences are included in our systematic errors. The solid lines in Figure 1 show the resulting fits and the Gaussian width parameters (σ) together with the integrated signal to background ratios ($\frac{S}{B}$) over the signal region ($\pi - 1 < \Delta\phi < \pi + 1$) are quoted.

The conditional yield (per trigger particle) is defined to be

$$CY = \frac{N_{pair}/\varepsilon_{assoc}}{N_{trig}}, \quad (3)$$

where ε_{assoc} ($\sim 0.15 \pm 0.015$) is the efficiency times acceptance for associated particles and N_{trig} is the number of trigger particles used to generate the correlation function. ε_{assoc} is obtained for each colliding system, centrality class, and transverse momentum bin by a GEANT based simulation of the PHENIX detector [14].




FIG. 2: Conditional yields are shown as a function of trigger particle pseudorapidity. The data points at mid-rapidity for $d + Au$ collisions are from [14]. To increase visibility, we artificially shift the data points belonging to the same η^{trig} bin. The errors on each point are statistical errors. The black bar around 0.1 on the left of the plot indicates a 10% common systematic error for all the data points due to the determination of associated particle efficiency. There is an additional $+0.037$ systematic error on the mid-rapidity $p + p$ point from jet yield extraction, which is shown as the arrow on that point (similar analysis as [16]).

It is interesting to plot the conditional yields as a function of η^{trig} . Changing η^{trig} from -2.0 to 2.0 effectively changes the range of Bjorken x of sampled partons in gold nuclei from $0.1^{+0.06}_{-0.04}$ to $0.01^{+0.02}_{-0.007}$. Results are shown in Figure 2. The first observation is that there is no difference beyond statistical fluctuations in the conditional yields for $p + p$, $d + Au$ peripheral, or $d + Au$ central collisions at any trigger particle pseudorapidity.

We further quantify any nuclear modification in the conditional yield by defining the following ratio:

$$I_{dAu} = \frac{CY|_{d+Au}}{CY|_{p+p}} \quad (4)$$

Figure 3 shows the ratio I_{dAu} as a function of p_T^{assoc} for central and peripheral $d + Au$ collisions, two different p_T^{trig} ranges, and for forward and backward rapidity trigger particles. In the plot, the shaded bands on each of the data points are the systematic errors which are the differences in N_{pair} obtained from the two methods described above and are independent from point-to-point. There is also a point-by-point correlated $\sim 2\%$ systematic uncertainty in the centrality dependence of ε_{assoc} determined by embedding Monte Carlo tracks into real events. The size of this uncertainty is comparable to the width of the $I_{dAu} = 1$ line.

Our measurement of the nuclear modification of the conditional yield indicates that I_{dAu} with the trigger particle at forward rapidity (sampling low- x partons in the gold nucleus) and backward rapidity (sampling high- x partons in the gold nucleus) are both consistent with one. There may even be some evidence of slight enhancement




FIG. 3: I_{dAu} vs. p_T^{assoc} for different centrality, p_T^{trig} and η^{trig} bins. To increase visibility, we artificially shift the data points belonging to the same p_T^{assoc} bin.




FIG. 4: The ratio of correlation peak widths between $d + Au$ and $p + p$ collisions. Only statistic error is shown. To increase visibility, we artificially shift the data points belonging to the same p_T^{assoc} bin.

for the case with trigger particles at forward pseudorapidity in central $d + Au$ collisions. We note that if mono-jets were a major contributor to our trigger particle sample in our x range, we would have expected a decrease in the conditional yield for $d + Au$ central collisions when the trigger particle is at forward pseudorapidity. Our measurement is inconsistent with any large nuclear suppression (*i.e.* mono-jets) of the jet structure in this kinematic range. However, we note that in these modest p_T ranges, there may be contributions from both “hard” (large Q^2) processes and “soft” coherent (small Q^2) processes. In $d + Au$ collisions “soft” particle production is shifted away from forward rapidity towards backward rapidity [21]. Thus, the fraction of hadrons at forward rapidity from “hard” processes may be increased in central $d + Au$ reactions, which may offer an explanation for the modest enhancement seen in the conditional yield for this case and could also mask off small mono-jet signals.

We have also compared the Gaussian widths of the

correlation peaks in $d + Au$ collisions to the widths in $p + p$ collisions. The ratios of the $d + Au$ widths to the $p + p$ widths are plotted in Figure 4 as a function of p_T^{assoc} . There may be a hint of a slight p_T^{assoc} dependence in the ratio, but overall there is no significant difference in the width in $d + Au$ collisions for different η^{trig} .

In conclusion, we have measured the two-particle azimuthal angle correlations in two centrality categories of $d + Au$ collisions and in $p + p$ collisions. We measured trigger particles at forward, backward and mid-rapidity and correlated them with associated particles at mid-rapidity. The associated particle conditional yields in central $d + Au$ collisions are consistent with the conditional yield in $p + p$ collisions. These conditional yields also do not change as we vary the trigger particle pseudorapidity over the range $|\eta| < 2.0$. We have also compared the widths of the away-side jet peaks in $d + Au$ and in $p + p$ collisions, and find no evidence for large η^{trig} -dependent modification.

We thank the staff of the Collider-Accelerator and Physics Departments at BNL for their vital contributions. We acknowledge support from the Department of Energy and NSF (U.S.A.), MEXT and JSPS (Japan), CNPq and FAPESP (Brazil), NSFC (China), IN2P3/CNRS, CEA, and ARMINES (France), BMBF, DAAD, and AvH (Germany), OTKA (Hungary), DAE and DST (India), ISF (Israel), KRF and CHEP (Korea), RMIST, RAS, and RMAE (Russia), VR and KAW (Sweden), U.S. CRDF for the FSU, US-Hungarian NSF-

OTKA-MTA, and US-Israel BSF.

* Deceased

† PHENIX Spokesperson:zajc@nevis.columbia.edu

- [1] I. Arsene *et al.*, Phys. Rev. Lett. **93**, 242303 (2004)
- [2] J. Adams *et al.*, Phys. Rev. **C 70**, 064907 (2004)
- [3] S. S. Adler *et al.*, Phys. Rev. Lett. **94**, 082302 (2005)
- [4] S. S. Adler *et al.*, Phys. Rev. Lett. **91**, 072303 (2003)
- [5] J. Adams *et al.*, Phys. Rev. Lett. **91**, 072304 (2003)
- [6] B. B. Back *et al.*, Phys. Rev. Lett. **91**, 072302 (2003)
- [7] I. Arsene *et al.*, Phys. Rev. Lett. **91**, 072305 (2003)
- [8] L. McLerran and R. Venugopalan, Phys. Rev. **D 49**, 2233 (1994); Phys. Rev. **D 49**, 3352 (1994)
- [9] R. Vogt, Phys. Rev. **C 70**, 064902 (2004)
- [10] J. Qiu, I. Vitev, hep-ph/0410218
- [11] R. C. Hwa, C. B. Yang and R. J. Fries, Phys. Rev. **C 71**, 024902 (2005)
- [12] D. Kharzeev, E. Levin and L. McLerran, Nucl. Phys. **A748**, 627 (2005)
- [13] Private communication with Ivan Vitev.
- [14] S. S. Adler *et al.*, nucl-ex/0510021
- [15] C. Adler *et al.*, Phys. Rev. Lett. **90**, 082302 (2005)
- [16] S. S. Adler *et al.*, Phys. Rev. **C 71**, 051902(R) (2005)
- [17] J. Adams *et al.*, Phys. Rev. Lett. **95**, 152301 (2005)
- [18] S. S. Adler *et al.*, nucl-ex/0507004
- [19] M. Arneodo, Phys. Rep. **240**, 301 (1994).
- [20] K. Adcox *et al.*, Nucl. Instrum. Methods **A499**, 469 (2003)
- [21] B. B. Back *et al.*, Phys. Rev. **C 72**, 031901 (2005)